


2016 ANNUAL REPORT

OF THE WATER QUALITY MONITORING PROJECT

FOR THE WATER QUALITY PROTECTION PROGRAM

OF THE FLORIDA KEYS NATIONAL MARINE SANCTUARY



Principal Investigators:
Henry O. Briceño¹ & Joseph N. Boyer²

¹Southeast Environmental Research Center
OE-148, Florida International University
Miami, FL 33199. <http://serc.fiu.edu/wqmnetwork/>

²Center for the Environment, Plymouth State University,
MSC 63, 17 High St., Plymouth, NH 03264

US EPA Agreement #X7-00049716-0

This is contribution number 842T from the Southeast Environmental Research Center in the Institute of
Water & Environment at Florida International University
June 2017

This page is intentionally left blank

2016 ANNUAL REPORT OF THE WATER QUALITY MONITORING PROJECT FOR THE WATER QUALITY PROTECTION PROGRAM OF THE FLORIDA KEYS NATIONAL MARINE SANCTUARY

Henry O. Briceño & Joseph N. Boyer

Funded by the Environmental Protection Agency (#X7-00049716-0)

EXECUTIVE SUMMARY

This report serves as a summary of our efforts to date in the execution of the Water Quality Monitoring Project for the FKNMS as part of the Water Quality Protection Program. The period of record for this report is Apr. 1995 – Dec. 2016 and includes data from 88 quarterly sampling events within the FKNMS (22 years).

Field parameters measured at each station (surface and bottom at most sites) include salinity (practical salinity scale), temperature ($^{\circ}\text{C}$), dissolved oxygen (DO, mg l^{-1}), turbidity (NTU), relative fluorescence, and light attenuation (K_d , m^{-1}). Water quality variables include the dissolved nutrients nitrate (NO_3^-), nitrite (NO_2^-), ammonium (NH_4^+), and soluble reactive phosphorus (SRP). Total unfiltered concentrations include those of nitrogen (TN), organic carbon (TOC), phosphorus (TP), silicate (SiO_2) and chlorophyll *a* (CHLA, $\mu\text{g l}^{-1}$).

The EPA developed Strategic Targets for the Water Quality Monitoring Project (SP-47) which state that beginning in 2008 through 2016, they shall annually maintain the overall water quality of the near shore and coastal waters of the FKNMS according to 2005 baseline. For reef sites, chlorophyll *a* should be less than or equal to $0.35 \mu\text{g l}^{-1}$ and the vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) should be less than or equal to 0.20 m^{-1} . For all monitoring sites in FKNMS, dissolved inorganic nitrogen should be less than or equal to $0.75 \mu\text{M}$ (0.010 ppm) and total phosphorus should be less than or equal to $0.25 \mu\text{M}$ (0.0077 ppm). Table 1 shows the number of sites and percentage of total sites exceeding these Strategic Targets for 2016.

We must recognize that the reduction of sampling sites in western FKNMS (less human-impacted sites) and the increase in inshore sites (heavily human-impacted sites) introduces a bias to the dataset which results in a reporting problem, perhaps requiring a revision of SP-47 to correct this deviation. To avoid such complications, we have not included the recently added locations (#500 to #509) in the calculation of compliances.

Table 1: EPA WQPP Water Quality Targets derived from 1995-2005 Baseline

For reef stations, chlorophyll less than or equal to 0.35 micrograms liter⁻¹ (ug l⁻¹) and vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) less than or equal to 0.20 per meter. For all stations in the FKNMS, dissolved inorganic nitrogen less than or equal to 0.75 micromolar and total phosphorus less than or equal to 0.25 micromolar. Water quality within these limits is considered essential to promote coral growth and overall health. The number of samples and percentage exceeding these targets is tracked and reported annually. Values in **green** are those years with % compliance greater than 1995-2005 **baseline**. Values in **yellow** are those years with % compliance less than 1995-2005 **baseline**.

EPA WQPP Water Quality Targets				
	REEF Stations		All Stations (excluding SHORE sites)	
Year	CHLA $\leq 0.35 \mu\text{g l}^{-1}$	$K_d \leq 0.20 \text{ m}^{-1}$	DIN $\leq 0.75 \mu\text{M}$	TP $\leq 0.25 \mu\text{M}$
			(0.010 mg l ⁻¹)	(0.008 mg l ⁻¹)
1995-05	1778 of 2367 (75.1%)	1042 of 1597 (65.2%)	7826 of 10254 (76.3%)	7810 of 10267 (76.1%)
2006	196 of 225 (87.1%)	199 of 225 (88.4%)	432 of 990 (43.6%)	316 of 995 (31.8%)
2007	198 of 226 (87.6%)	202 of 222 (91.0%)	549 of 993 (55.3%)	635 of 972 (65.3%)
2008	177 of 228 (77.6%)	181 of 218 (83.0%)	836 of 1,000 (83.6%)	697 of 1,004 (69.4%)
2009	208 of 228 (91.2%)	189 of 219 (86.3%)	858 of 1,003 (85.5%)	869 of 1,004 (86.6%)
2010	170 of 227 (74.9%)	176 of 206 (85.4%)	843 of 1,000 (84.3%)	738 of 1,003 (73.6%)
2011	146 of 215 (67.9%)	156 of 213 (73.2%)	813 of 1,012 (80.3 %)	911 of 1,013 (89.9 %)
2012	142 of 168 (84.5%)	135 of 168 (80.4%)	489 of 683 (71.6 %)	634 of 684 (92.7 %)
2013	148 of 172 (86.0%)	150 of 172 (87.2%)	496 of 688 (72.1 %)	603 of 688 (87.6 %)
2014	141 of 172 (82.0%)	133 of 172 (77.3%)	426 of 690 (61.7%)	540 of 690 (78.3%)
2015	122 of 172 (70.9%)	135 of 172 (78.5%)	487 of 688 (70.8%)	613 of 688 (89.1%)
2016	131 of 172 (76.2%)	129 of 170 (75.9%)	427 of 687 (62.2%)	549 of 688 (79.8%)

Important Results Realized from this Monitoring Project

1. Land-based Influence on Water Quality

We documented elevated nutrient concentrations (DIN, TN, TP, and SiO₂) in waters close to shore along the Keys, and their corresponding responses from the system, such as higher phytoplankton biomass (CHLA), turbidity, as well as lower salinity and DO in the water column. These changes, associated to human impact, have become even more obvious by the addition of 10 stations located close to shore, (SHORE, within the 500 m Halo; FDEP 2011) sampled since Nov 2011 (Fig i).




Figure i. Nutrient and response changes along transect from shore sites to reef-track

This trend, especially for DIN was evident from our first sampling event in 1995 and was not observed in a comparison transect from the Tortugas (no human impact). This pattern suggests a land-

based, freshwater nutrient source. In summary, this type of distribution would imply a relatively elevated land source which is diluted by low nutrient Atlantic Ocean waters. However, this is not always the case as deep water upwelling to the reef occurs on a regular basis

2. Trend analysis – 22 years

Surface DO has increased in most areas of the FKNMS (Fig. ii). Greatest increases in surface DO occur on the Atlantic side of the Keys and in some inshore areas on the Bay side. Bottom DO trends showed a similar pattern as surface with more increased DO than surface sites (Fig. iii). Increased DO is beneficial for animal life.




Figure ii. Total change in DO of surface waters for 22 year period calculated from significant trends.




Figure iii. Total change in DO of bottom waters for 22 year period calculated from significant trends.

Water column turbidity (cloudiness) declined throughout the FKNMS during the 22 year period (this is a good thing). There was no significant change in turbidity in bottom waters. The largest declines in turbidity occurred in western Florida Bay and west of the Marquesas (Fig iv).




Figure iv. Total change in Turbidity in surface waters for 22 year period calculated from significant trends.

Although light extinction (K_d) did not show significant trend, decreased turbidity did affect the percent of surface light (I_o) reaching the bottom. Bottom light increased at most reef sites throughout the Keys (Fig. v). More light on the bottom is beneficial to corals, seagrass, and algae. Interestingly, the Backcountry area of the lower Keys experienced increases in K_d which lead to corresponding decreases in bottom light.




Figure v. Total change in bottom I_o for 22 year period calculated from significant trends.

No significant trends in TP were detected, however very small increases in SRP, up to 1.5 ppb over 22 years, were observed for most surface waters (Fig. vi). Concentrations of SRP are generally an order of magnitude lower than TP.




Figure vi. Total change in SRP in surface waters for 22 year period calculated from significant trends.

The largest sustained monotonic trend has been the decline in surface TOC, especially in the Backcountry and the Marquesas (Fig. vii). This is part of a regional trend in TOC observed on the SW Shelf, Florida Bay, and the Everglades mangrove estuaries. This decline could be considered favorable given that TOC is an important component of water color and affects light penetration, but could also be an indication of decreased upstream primary production.




Figure vii. Total change in TOC in surface waters for 22 year period calculated from significant trends.

The large scale of this monitoring program has allowed us to assemble a much more holistic view of broad physical/chemical/biological interactions occurring over the South Florida hydroscope. This confirms that rather than thinking of water quality monitoring as being a static, non-scientific pursuit it should be viewed as a tool for answering management questions and developing new scientific hypotheses. We continue to maintain a website (<http://serc.fiu.edu/wqmnetwork/>) where data and reports from the FKNMS are integrated with other programs.